



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> <b>C07K 7/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/75173</b> <b>(43) International Publication Date:</b> 14 December 2000 (14.12.2000)
<b>(21) International Application Number:</b> PCT/GB00/02128 <b>(22) International Filing Date:</b> 02 June 2000 (02.06.2000) <b>(30) Priority Data:</b> 9912852.2 02 June 1999 (02.06.1999) GB <b>(60) Parent Application or Grant</b> REGEN THERAPEUTICS PLC [/]; (). GEORGIADES, Jerzy, A. [/]; (). GEORGIADES, Jerzy, A. [/]; (). CURTIS, Philip, Anthony; ().		<b>Published</b>
<b>(54) Title: PEPTIDES</b> <b>(54) Titre: PEPTIDES</b>  <b>(57) Abstract</b> <p>The amino acid sequence of several peptides present in Colostrinin is disclosed. These peptides are useful, inter alia, in the treatment of disorders of the immune system and the central nervous system.</p> <b>(57) Abrégé</b> <p>L'invention concerne la séquence d'acides aminés de plusieurs peptides présents dans la Colostrinine. Lesdits peptides sont utiles, entre autres, dans le traitement de troubles du système immunitaire et du système nerveux central.</p>		

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



RECEIVED  
PCT

(43) International Publication Date  
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number  
WO 00/75173 A2

- (51) International Patent Classification<sup>7</sup>: C07K 7/00 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (21) International Application Number: PCT/GB00/02128
- (22) International Filing Date: 2 June 2000 (02.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9912852.2 2 June 1999 (02.06.1999) GB
- (71) Applicant (*for all designated States except US*): REGEN THERAPEUTICS PLC [GB/GB]; 88 Kingsway, London WC2B 6AA (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (*for US only*): GEORGIADES, Jerzy, A. [US/US]; 9615 Bayou Brook, Houston, TX 77063 (US).
- (74) Agents: CURTIS, Philip, Anthony et al.; A.A. Thornton & Co., 235 High Holborn, London WC1V 7LE (GB).
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— Without international search report and to be republished upon receipt of that report.

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 00/75173 A2

(54) Title: PEPTIDES

(57) Abstract: The amino acid sequence of several peptides present in Colostrinin is disclosed. These peptides are useful, inter alia, in the treatment of disorders of the immune system and the central nervous system.

PEPTIDES

The present invention relates to peptides. More particularly the invention relates to certain peptides isolated from Colostrinin. The invention also relates to therapeutic uses of the peptides and to antibodies derived therefrom.

Colostrum is the thick, yellowish fluid produced by a mammalian mother's breasts during the first few days after childbirth. It is the first lacteal secretion post parturition and it contains a high concentration of immunoglobulins (IgG, IgM and IgA) and other proteins. It is replaced by mature breast milk about four to five days after birth. Compared with mature breast milk, colostrum contains low sugar and iron, but is rich in lipids, proteins, mineral salts, vitamins and immunoglobulins. Colostrum also contains various floating cells such as granular and stromal cells, neutrophils, monocyte/macrophages and lymphocytes and includes growth factors, hormones, cytokines and polypeptide complexes.

Various factors have been isolated and characterised from mammalian colostrum. In 1974, Janusz et al (FEBS Lett., 49, 276-279) isolated a proline-rich polypeptide (PRP) from ovine colostrum. It has since been discovered that mammals other than sheep have analogues of PRP as a component of their colostrum. PRP has since been called Colostrinin (and is sometimes called Colostrinine).

M. Janusz & J. Lisowski in "Proline-Rich Polypeptide (PRP) - an Immunomodulatory Peptide from Ovine Colostrum" (Archivum Immunologiae et Therapiae Experimentalis, 1993, 41, 275-279) mentioned that PRP from ovine colostrum has immunotropic activity in mice.

A. Dubowska-Inglot et al in "Colostrinine: a proline-rich polypeptide from ovine colostrum is a modest cytokine inducer in human leukocytes" (Archivum Immunologiae et Therapiae Experimentalis, 1996, 44, 215-224) discussed the use of Colostrinin in the treatment of Alzheimer's disease. The use of Colostrinin in the treatment of Alzheimer's disease, and other conditions, was also discussed in WO-A-98/14473 and in "Colostrinin: a Proline-Rich Polypeptide (PRP) Complex Isolated from Ovine Colostrum for Treatment of Alzheimer's Disease. A Double-Blind, Placebo-Controlled Study", Leszek, J. et al, Archivum Immunologiae et Therapiae Experimentalis, 1999, 47, 377-

385.

Colostrinin, in its natural form, is obtained from mammalian colostrum. As described in WO-A-98/14473, analysis by electrophoresis and chromatography has shown that Colostrinin has the following properties:

- (i) it has a molecular weight in the range 16,000 to 26,000 Daltons (this was shown by electrophoresis in the presence of SDS);
- (ii) it is a dimer or trimer of sub-units each sub-unit having a molecular weight in the range 5,000 to 10,000 Daltons (this was shown by acrylamide gel electrophoresis in the presence of SDS);
- (iii) it contains proline, and the amount of proline is greater than the amount of any other single amino acid (this can be shown by conventional amino acid analysis).

By means of these techniques it was shown that ovine Colostrinin has a molecular weight of about 18,000 Daltons, is made up of three non-covalently linked sub-units each having a molecular weight of about 6,000 Daltons and includes about 22 wt% proline. The amino-acid composition of ovine Colostrinin was shown to be made up of the following number of residues per sub-unit: lysine - 2, histidine - 1, arginine - 0, aspartic acid - 2, threonine - 4, serine - 3, glutamic acid - 6, proline - 11, glycine - 2, alanine - 0, valine - 5, methionine - 2, isoleucine - 2, leucine - 6, tyrosine - 1, phenylalanine - 3 and cysteine - 0.

We have now further analysed the composition of Colostrinin in order to try to identify its components, so that a synthetic form of Colostrinin can be produced.

We have concluded that Colostrinin contains peptide fragments from at least two different proteins: annexin; and  $\beta$ -casein. In addition, Colostrinin contains a number of other peptide fragments which do not have any known precursor protein; these amino acid sequences may be derived from an unknown precursor protein, or they may have no precursor protein. It is believed that some of the peptide sequences are from a  $\beta$ -casein homologue.

According to one aspect of the present invention there is provided a peptide having one of the following amino acid sequences A-1 to D-1:

-3-

5

*Group A: Peptides of unknown precursor*

A-1 LQTPQPLLQVMMEPQGD

10

A-2 MPQNFYKLPQM

A-3 VLEMKFPPPPQETVT

5

A-4 LKPFPKLVVEVFPFP

A-5 SEQP

15

A-6 DKE

A-7 DPPPQS

A-8 LNF

20

10

*Group B: Peptides (possibly) having  $\beta$ -casein homologue precursor*

B-1 VLPPNVG

B-2 KYKLQPE

B-3 SEEMP

25

B-4 DSQPPV

15

B-5 FPPPK

B-6 VVMEV

30

B-7 DLEMPVLPVEFPFV

B-8 LFFFLPVVNVLP

B-9 MQPPPLP

20

B-10 DQPPDVEKPDLPFQVQS

35

*Group C: Peptides having  $\beta$ -casein precursor*

C-1 VYPFTGIPN (Casein Position 74-83)

C-2 SLPQNILPL (Casein Position 84-92)

40

C-3 TQTPVVVPPF (Casein Position 93-102)

25

C-4 LQPEIMGVPKVKETMVPK (Casein Position 103-120)

C-5 HKEMPFKYPVEPFTESEQ (Casein Position 121-138)

C-6 SLTLTDVEKLHLPLLVQ (Casein Position 139-156)

45

C-7 SWMHQPP (Casein Position 157-163)

C-8 QPLPPTVMFP (Casein Position 164-173)

30

C-9 MHQPPQPLPPTVMFP (casein Position 159-173)

50

C-10 PQSVLS (Casein Position 174-179)

55

-4-

C-11 LSQPKVLPVPQKAVPQRDMPIQ (Casein Position 180-201)

C-12 AFLLYQE (Casein Position 202-208)

C-13 FLLYQEPVLGPVR (Casein Position 203-214)

C-14 RGPFPILV (Casein Position 214-222)

5 *Group D: Peptides having annexin precursor*

D-1 ATFNRYQDDHGEEILKSL (Annexin Position 203-220)

15 It is possible that the peptides in group A are also derived from the  $\beta$ -casein homologue, but there is currently no evidence to support this conclusion.

20 These peptides may be provided in substantially isolated form. Furthermore, a composition may be provided which contains two or more of the above peptides, in combination.

25 In respect of the peptides A-1 to B-10, the invention further includes any peptide which includes the specified amino acid sequence. In respect of the peptides A1 to D1, the invention further comprises any peptide which includes an amino-terminal amino acid sequence corresponding to the specified sequence. Thus, with reference to peptide A-1, for example, the invention encompasses any peptide having the N-terminal amino acid sequence LQTPQPLLQVMMEPQGD; the same applies to peptides A-2 to D-1. For the avoidance of doubt, it is stated that the amino-terminal end is on the left hand side of the sequence, in accordance with the usual convention. It will be appreciated that any of the specified amino acid sequences may be provided with an inert amino acid sequence on the amino-terminal and/or the carboxy-terminal end thereof. The invention further includes physiologically acceptable active derivatives of the peptides.

40 The peptides can be obtained by a number of techniques. In one embodiment, they can be prepared naturally by isolation from Colostrinin or colostrum. In a preferred embodiment, they are prepared by a conventional technique for peptide synthesis, such as by solid-phase or liquid-phase peptide synthesis. Alternatively, the gene sequence encoding the peptides can be constructed by known techniques such as expression vectors or plasmids and transfected into suitable microorganisms that will express the DNA sequences, whereby the peptides can be later extracted from the medium in which the microorganisms are grown. Thus, the invention also embraces a DNA sequence

5 encoding the peptides described above, and a recombinant vector prepared by inserting said DNA in a vector.

10 The peptides, either alone or in combination with one another, have a number of therapeutic uses.

5 In one advantageous embodiment, one or more of peptides A-1 to D-1 may be used in the treatment of disorders of the central nervous system, particularly chronic  
15 disorders of the central nervous system. The disorders of the central nervous system that may be treated include neurological disorders and mental disorders. Examples of neurological disorders that may, with advantage, be treated include dementia, and also  
20 disorders that cause dementia, such as neurodegenerative disorders. Neurodegenerative disorders include, for example, senile dementia and motor neurone disease; Parkinson's disease is an example of a motor neurone disease that can be treated. Alzheimer's disease is an example of a neurodegenerative disease that can  
25 be treated. Examples of mental disorders that can be treated by one or more of the 15 peptides include psychosis and neurosis. For example, the peptides may be used to treat emotional disturbances, especially the emotional disturbances of psychiatric patients in a state of depression. The peptides may also be used as an auxiliary withdrawal treatment for drug addicts, after a period of detoxification, and in persons dependent on stimulants.

20 In another advantageous embodiment of the invention, one or more of peptides A-1 to D-1 may be used in the treatment of disorders of the immune system, particularly chronic disorders of the immune system the may occur spontaneously in  
35 people of advanced age. The peptides can also be used in the treatment of diseases requiring immuno-modulation. The peptides are useful in the treatment of a variety of  
40 25 diseases with an immunological and infectious basis. For example, they can be used to treat chronic diseases with a bacterial and viral aetiology, and to treat acquired immunological deficiencies that have developed, for example, after chemotherapy or radiotherapy of neoplasms. The peptides may be used for treating chronic bacterial and viral infections requiring non-specific immunostimulation and immunocorrection.

30 A chronic disorder is a disorder that has persisted, or is expected to persist, for a long time, i.e., at least 3 months and usually at least 6 months.

5 One or more of the peptides may be used for improving the development of the  
immune system of a new born child. It is a further feature of the invention to use the  
10 peptides to correct immunological deficiencies in a child. These uses of the peptides  
may be particularly applicable to babies or children who have been deprived of  
5 colostrum. This may occur, for example, in babies and children who were not breast fed  
from birth.

15 The peptides, either alone or in combination with one another, also have  
diagnostic and research applications. For example, the synthetic peptides, as well as  
the corresponding antibodies described below, may be used to recognise pathological  
20 10 processes occurring in a host. These processes may be induced by excessive  
production or inhibition of the peptides or the antibodies. Once the pathological process  
associated with a particular level of the peptides or the antibodies is known, measuring  
the production of the peptides and the antibodies in body fluids may be used to  
25 determine pathological processes taking place in the host.

15 According to another aspect of the invention, we provide the use of one or more  
of peptides A-1 to D-1 as a dietary supplement. This dietary supplement is particularly  
30 useful for babies, especially premature babies and babies at term, and for young  
children to correct deficiencies in the development of their immune system. The dietary  
supplement may also be used as a dietary supplement for adults, including senile  
20 persons, who have been subjected to chemotherapy, or have suffered from anorexia,  
or weight loss due to chronic disease.

35 In an aspect of the invention, we provide a dietary supplement comprising an  
orally ingestible combination of one or more of peptides A-1 to D-1 in combination with  
a physiologically acceptable carrier. The dietary supplement may be provided in liquid  
40 25 or solid form; the dietary supplement may suitably be provided in the form of a tablet.  
The dietary supplement may be provided in the form of a baby food formula. The  
dietary supplement may include, as an additive, lactoferrin and/or selenium and/or a  
45 group of cytokines containing members of the interferon family.

In accordance with the invention, one or more of peptides A-1 to D-1 may be  
30 administered prophylactically in order to help to prevent the development of disorders  
of the central nervous system and the immune system.



5 The peptides according to the invention may be used to promote the dissolution  
of  $\beta$ -amyloid plaques, and, therefore, the peptides may be used in the treatment of any  
10 disease which is characterised by the development of  $\beta$ -amyloid plaques.

The peptides according to the invention may be administered in a dosage in the  
5 range 1 ng to 10 mg. A dosage unit of about 3  $\mu$ g is typical. However, the optimum  
dosage will, of course, depend upon the condition being treated.

15 The peptides according to the invention may be formulated for administration in  
any suitable form. Thus, the invention further provides a composition, especially a  
pharmaceutical composition, which includes one or more of the peptides in combination  
20 with a physiologically acceptable carrier. The peptides may, for example, be formulated  
for oral, topical, rectal or parenteral administration. More specifically, the peptides may  
be formulated for administration by injection, or, preferably, in a form suitable for  
absorption through the mucosa of the oral/nasopharyngeal cavity, the alimentary canal  
25 or any other mucosal surface. The peptides may be formulated for administration  
15 intravenously, subcutaneously, or intramuscularly. The oral formulations may be  
provided in a form for swallowing or, preferably, in a form for dissolving in the saliva,  
whereby the formulation can be absorbed in the mucous membranes of the  
30 oral/nasopharyngeal cavity. The oral formulations may be in the form of a tablet for oral  
administration, lozenges (i.e. a sweet-like tablet in a form suitable to be retained in the  
20 mouth and sucked), or adhesive gels for rubbing into the gum. The peptides may be  
35 formulated as an adhesive plaster or patch, which may be applied to the gums. The  
peptides may also be formulated for application to mucous-membranes of the genito-  
urinary organs. The topical formulations may be provided in the form of, for example,  
40 a cream or a gel.

25 One or more of the peptides may be incorporated into products like milk or  
cheese spread.

45 According to another aspect of the invention there is provided a pharmaceutical  
composition comprising a peptide containing the amino acid chain  
LQTPQPLLQVMMEPQGD; DPPPPQS; and/or LFFFLPVVNVLP or use as an  
30 immunosuppressant, for use in the treatment of autoimmune disorder, and/or for use  
50 in suppressing the rejection of transplanting organs. The invention also embraces the

5 use of one or more of these peptides in the manufacture of a medicament for use as an immunosuppressant, for use in the treatment of autoimmune disorder, and/or for use  
10 in suppressing the rejection of transplanting organs.

We have found that the ratio of the peptides in colostrum varies over time.  
5 Owing to hormonal changes, many proteins secreted into colostrum become sequentially degraded. The longer the time from parturition the more extensive the  
15 degradation can be. This knowledge will help with the design of new baby food formulas as well as many drugs for immuno-compromised patients.

In another aspect, the invention provides an antibody for each of the peptides  
20 A-1 to D-1, and provides compositions containing said antibodies. In particular the invention provides the antibodies in substantially isolated form. The antibodies can be produced by injecting a suitable mammalian subject, such as a rabbit, with the corresponding peptide (with a suitable adjuvant), then recovering the antibodies from  
25 the subject after allowing time for them to be produced. This technique is described in detail in Example 3. It is possible to test that the correct antibody has been produced  
15 by ELISA (enzyme-linked immunosorbent assay) using the synthetic peptides as antigens. The antibodies can be further tested against the natural peptides in Colostrinin as confirmation that the synthetic peptides do correspond to the natural peptides found in Colostrinin. The antibodies have potential uses in therapy, as a  
30 diagnostic tool and as a research tool.

35 The invention also encompasses the selective administration of one or more of peptides A-1 to D-1, at selected times to a patient, and the selective administration of one or more of the antibodies for the peptides in order to switch on or off the activity of the peptides at a selected time.

40 25 A selection of selected ones of the peptides and/or antibodies may be provided in a single composition which is specially tailored to produce a particular effect. For example, for a person with an immunological disorder, the composition can be specially  
45 tailored for that disorder. The composition may be specially selected for more than one disorder. The composition may be specially selected to restore or produce a particular  
30 balance in a subject.

50 In some applications it may be desirable to provide a pharmaceutical

composition which contains one or more of the peptides and one or more of the antibodies in combination with a physiologically acceptable carrier.

The invention further embraces the use of one or more of the peptides and/or antibodies in the manufacture of a medicament for use in any of the therapeutic applications described above.

Reference is now made to the accompanying drawing in which Figs. 1 to 18 are Matrix-Assisted Laser Desorption Time-of-Flight Mass Spectroscopy (LDMS) spectra of certain peptides according to the invention.

The invention will now be further described with reference to the following examples.

#### Example 1

##### Preparation of Colostrinin

Colostrinin can be prepared by techniques already disclosed in the prior art, including, for example, WO-A-98/14473. Colostrum collected from the ewe within 12 hours post parturition can be purified by centrifuging to eliminate cellular and lipidic components, pH shifting to eliminate nutritional components, ammonium sulfate precipitation, ion exchange chromatography and molecular sieving.

#### Example 2

##### Identification of the Components of Colostrinin

Initially the Colostrinin produced according to example 1 was analysed by SDSPAGE, by means of which we found the following two peptides: VLEMKFPPPPQETVT (A-3) and LKPFKLKVEVFPEP (A-4). However, we could not identify any other peptides with this technique, so we turned to hplc.

The Colostrinin produced in example 1 was fractionated by hplc using a C-18 reverse-phase column. This technique was used to separate the peptides exhibiting different hydrophobic patterns, present in Colostrinin. The hplc column was obtained from Separation Methods Technologies (who are based in Newark, Delaware, U.S.A). The column type was designated C-18 and was 150 mm in length by 10 mm in diameter. The column was packed with particles having a particle size of 3  $\mu$ m having

-10-

a pore size of 30 nm. The pump module and diode array were supplied by Beckman (who are based in Fullerton, California, U.S.A.): a Beckman System Gold 126 pump module was used, and a Beckman System Gold 168 diode array detector module was used.

The Colostrinin was loaded in 0.1% trifluoroacetic acid (TFA) dissolved in hplc grade water. A 500 µl sample, containing approximately 900 picomole of the Colostrinin was loaded on the column, the column having been equilibrated prior to loading. After approximately 10 minutes of intensive washing, the material was eluted by gradient formed from solutions A and B, under a regime indicated in Table 1. During this time, the flowrate through the column was 0.06 ml/min.

Table 1

Time/Min	% Solvent A	% Solvent B
0.00	95.0	5.0
10.00	30.0	70.0
100.00	0.0	100.0
140.00	95.0	5.0
150.00	95.9	5.0

Solvent A: 0.1% TFA (trifluoroacetic acid) in hplc grade water.

Solvent B: 70% acetonitrile fluoride and 0.09% TFA in hplc grade water.

The peptides found at the peaks in the hplc were then individually analysed using Edman Degradation; this was done using a Beckman LF3000 sequencer. Each concentrated fraction was loaded into a pre-salted Beckman peptide support disk. The samples were sequenced using the standard Edman degradation steps. Typically, 10 to 100 pmoles were used to generate 10 to 25 cycles for each analysis.

Subsequently, each fraction was analysed by the Inline hplc System. This used a Hewlett Packard PTH-AA column having a length of 250 mm and a diameter of 2.1 mm. The Beckman System Gold 126 pump module was used, and the Beckman System Gold 168 diode array detector module was used. The flowrate in the column was 0.275

5

-11-

ml/min, and the solvent composition was varied as shown in Table 2.

10

15

20

25

30

35

40

45

50

55

-12-

Table 2

Time/Min	% Solvent A	% Solvent B
0.00	80.0	20.0
0.10	62.0	38.0
17.10	10.0	90.0
28.10	87.5	12.5

Solvent A: 3.5% THF (tetrahydrofuran), 1.5% acetonitrile fluoride premix, 1% acetic acid & 0.02% TEA (triethanolamine) in hplc grade water.

Solvent B: 12% isopropanol in acetonitrile.

The structure of the peptides A-1 to D-1 was then used for comparative studies with sequences registered in two known computer programs: Wu-Blast 2 of the National Center for Biotechnology Information NR Protein Data Base; and Beauty - Post Processing provided by the Human Genome Center, Baylor College of Medicine, Houston, Texas, USA. This made it possible to determine whether any of the peptide sequences P1-P32 were already known.

The results of the Edman degradation are summarised in Table 3. The subsequent analysis with the computer programs revealed that there were at least two different precursor proteins for the peptides in Colostrinin:  $\beta$ -casein and annexin. Furthermore, by using the Tremble program, it was possible to find evidence that some of the peptides may have a precursor which is a casein homologue. Finally, some of the peptides had unique sequences with no homology to any known protein.

Table 3

Peak No.	Elution time min.	Area %	AA sequence		
			Casein homologue	Unknown precursor	Casein/Annexin precursor
1	8.54	1.181	VMEV (B-6)		ATFNRYQDDHGEEILKSL (D-1)

	2	29.086	0.124		SEQP (A-5)	
	3	53.775	0.579			
10	4	56.815	0.111	FPPPK (B-5)		LSQPKVLPVPQKAPQPRDM PIQ (C-11)
	5	58.044	2.101	DSQPPV (B-4)		LSQPKVLPVPQKAPQPRDM PIQ (C-11)
15	5	60.488	0.588	MQPPPLP (B-9)		LSQPKVLPVPQKAPQPRDM PIQ (C-11)
	7	62.684	1.273		DPPPPQS (A-7)	
20	8	65.44	3.247		LQTPQPLLQV MMEPQGD (A-1)	LSQPKVLPVPQKAPQPRDM PIQ (C-11)
	9	66.775	0.683	DQPPDVEKPDQ PFQVQS (B-10)		LSQPKVLPVPQKAPQPRDM PIQ (C-11)
25	10	67.929	2.943	LFFFLPVVNVLP (B-8)		LSQPKVLPVPQKAPQPRDM PIQ (C-11) MHQPPQPLPPTVMFP (C-9)
30	10	69.229	2.717	SEEMP (B-3)		LSQPKVLPVPQKAPQPRDM PIQ (C-11) HKEMPFKYVPEPFTESQ (C-5)
35	12	70.984	2.964	KYKLQPE (B-2)		LSQPKVLPVPQKAPQPRDM PIQ (C-11) HKEMPFKYVPEPFTESQ (C-5)
40	13	72.547	1.423	VLPPNVG (B-1)		LSQPKVLPVPQKAPQPRDM PIQ (C-11)
	14	74.09	1.425	DLEMPVLPVEPF PFV (B-7)		SLPQNILPL (C-2)
45	15	76.558	5.268		MPQNFYKLP QM (A-2)	MHQPPQPLPPTVMFP (C-9)
15	16	78.506	6.978		LNF (A-8)	MHQPPQPLPPTVMFP (C-9)

5

10

15

5

20

25

10

30

35

40

45

50

17	80.94	4.224			MHQPPQPLPPTVMFP (C-9) SLTLTDVEKLHLPLPLVQ (C-6) PQSVLS (C-9)
18	83.8	1.025			ND
19	84.314	2.151			MHQPPQPLPPTVMFP (C-9)
20	85.707	3.103			SWMHQPP (C7)
21	87.061	1.047			ND
22	87.907	1.529			ND
23	88.921	1.311			MHQPPQPLPPTVMFP (C-9) SLTLTDVEKLHLPLPLVQ (C-6) TQTPVVPPF (C-3) VYPFTGPIPN (C-1)
24	89.856	1.114			ND
25	91.343	0.906			ND
26	92.667	0.821			ND
27	93.521	3.893			ND
28	94.751	1.426			ND
29	95.82	0.272			HKEMPFKYPVEPFESQ (C-5)
30	96.697	3.164			QPLPPTVMFP (C-8) HKEMPFKYPVEPFESQ (C-5)
31	97.938	3.266			ND
32	99.893	5.621			HKEMPFKYPVEPFESQ (C-5)
33	100.9	5.032			ND
34	102.709	4.007			AFLLYQE (C-12) HKEMPFKYPVEPFESQ (C-5)
35	104.74	3.275			ND

55



36	106.01	2.231			ND
37	170.75	3.037			ND
38	108.782	2.173			SLTLTDVEKLHLPLPLVQ (C-6) HKEMPFPKYPVEPFOTESQ (C-5) SLPQNILPL (C-2) VYPFTGPIPN (C-1)
39	111.056	5.375			HKEMPFPKYPVEPFOTESQ (C-5)
40	112.679	1.901			ND
41	114.707	0.436			ND
42	8.54	1.181			ATFNRYQDDHGEEILKSL (D-1)

ND indicates that these fractions were not analysed.

DKE (A-6), LQPEIMGVVKVETMVPK (C-4), FLLYQEPVLGPVR (C-11) and RGPFPILV (C-13) were also detected by hplc, although their presence is not indicated in the above table.

### Example 3

#### Production of the Antibodies

The peptides identified in example 2 were produced by the synthetic technique known as the solid phase method. This method involved the following steps:

1. Wash pre-loaded resin with DMF (dimethylformamide), then drain completely.
2. Add 10 ml of 20% piperidine/DMF to resin. Shake for 5 mins, then drain.
3. Add another 10 ml of 20% piperidine/DMF. Shake for 30 mins.
4. Drain reaction vessel and wash resin with DMF four times. Then wash once with DCM (dichloromethanol). Check beads using the ninhydrin test - the beads should be blue.
5. The coupling step was carried out as follows:

Prepare the following solution:

1 mmole Fmoc (i.e. fluorenylmethyloxycarbonyl) amino acid

-16-

2.1 ml of 0.45 M HBTU/HOBT (1 mmol) (2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate/N-hydroxybenzotriazole-H<sub>2</sub>O)

348 µl of DIEA (2 mmol) (diisopropylethylamine)

Add the solution to the resin and shake for a minimum of 30 minutes.

6. Drain reaction vessel and wash the resin again with DMF four times and with DCM once.

7. Perform the ninhydrin test:

If positive (no colour) - proceed to step 2 and continue synthesis.

If negative (blue colour) - return to step 5 and recouple the same Fmoc amino acid.

8. After the synthesis was complete, the peptide was cleaved from the resin with 5% H<sub>2</sub>O, 5% phenol, 3% Thionisole, 3% EDT (ethanedithiol), 3% triisopropylsilane and 81% TFA for 2 hours.

9. After 2 hours, filter into cold MTBE (methyl t-butyl ether). The precipitated peptide was then washed twice with cold MTBE and dried under nitrogen gas.

10. The molecular weight of the synthesised peptides was checked by Matrix-Assisted Laser Desorption Time-of-Flight Mass Spectroscopy (LDMS), and the purity was checked by hplc using a C-18, 300 Angstrom, 5 µm column. The resulting spectra of some peptides are shown in Figs. 1 to 18.

To each N-terminal end of the synthetic peptides, L-cysteine was attached, and the peptide was formed into a ring so that the cysteine group lay between the N-terminal and the C-terminal ends of the synthetic peptide. This facilitated peptide conjugation with Keyhole Hemolymph (KHL). The shorter peptides (i.e. those containing 9 or fewer amino acids) were artificially elongated with biologically inert amino acids prior to attaching the L-cysteine. This was done in order to facilitate annealing and increase the antigenicity of the shorter peptides.

Table 4 shows a number of the peptides that were formed and indicates the figure

number of the drawings which illustrates the laser desorption mass spectrum.

Table 4

5	PEPTIDE SYNTHESISED	ORIGINAL PEPTIDE	FIGURE NO.
15	NH <sub>2</sub> -(Ac)CLQTPQPLLQVMMEPQGD-OH	A-1	1
	NH <sub>2</sub> -(Ac)CMPQNFYKLPQM-OH	A-2	2
	NH <sub>2</sub> -(Ac)CVLEMKFPPPPQETVT-OH	A-3	3
20	NH <sub>2</sub> -(Ac)CLKPFPKLKVEVFPFP-OH	A-4	4
10	NH <sub>2</sub> -SEQPGGGC-OH	A-5	5
	NH <sub>2</sub> -(Ac)CGVLPPNVG-OH	B-1	6
	NH <sub>2</sub> -(Ac)CGGGKYKLQE-OH	B-2	7
25	NH <sub>2</sub> -(Ac)CGGGSEEMP(amide)-OH	B-3	8
	NH <sub>2</sub> -(Ac)CGGGDSQPPV-OH	B-4	9
15	NH <sub>2</sub> -CFPPPKGGGC-OH	B-5	10
30	NH <sub>2</sub> -(Ac)CGGGVMEV-OH	B-6	11
	NH <sub>2</sub> -(Ac)CDLEMPVLPVEFPFV-OH	B-7	12
	NH <sub>2</sub> -(Ac)CLFFFLPVVNVLP-OH	B-8	13
35	NH <sub>2</sub> -(Ac)CMQPPPLP-OH	B-9	14
20	NH <sub>2</sub> -(Ac)CDQPPDVEKPDLPFQVQS-OH	B-10	15
	NH <sub>2</sub> -(Ac)CGAFLLYQE-OH	C-12	16
	NH <sub>2</sub> -(Ac)CATFNRYQDDHGEEILKSL-OH	D-1	17
40	NH <sub>2</sub> -DPPQSGGGC-OH	A-7	18

25 The invention further provides each of the peptides specified in Table 4, and the  
 45 cyclised version of each of these peptides, especially in isolated form and produced  
 by a synthetic process. The term "Ac" represents an acyl group.

For immunisation, two young adult rabbits (5-6 months old, weighing 5-6 lbs [2.3-  
 2.7kg]) were used. Each antigen (i.e., each synthetic peptide) was given subcutaneously  
 50 and intramuscularly in 0.1 ml injections at ten different sites. The protocol used followed

the following sequence:

	<u>Day</u>	<u>Procedure</u>
5	0	Prebleed & initial inoculation of rabbit with 200 µg of the peptide at
10	5	0.5 ml of conjugate solution mixed with an equal volume of
15		complete Freund's adjuvant (mineral oil/emulsifier/killed
	14	mycobacteria).
20	10	Boost inoculation with 200 µg of the peptide at 0.5 ml of conjugate
	28	solution mixed with an equal volume of incomplete Freund's
		adjuvant (mineral oil/emulsifier).
25		Boost (as on day 14)
	42	Production Bleed (approx. 20ml sera)
		Boost (as on day 14)
		Production Bleed (approx. 20ml sera)
30	15	Boost (as on day 14)
	56	Production Bleed (approx. 20ml sera)
	70	Boost (as on day 14)
		Production Bleed (approx. 20ml sera)

20 This protocol may be varied. For example, the frequency of the production bleed  
35 depends upon, inter alia, the size and health of the host species.

The sera produced by this protocol were used for IgG purification on a Protein  
A matrix (from Sigma, based in St. Louis, MO, USA). The protocol was as follows:

- 40 1. Wash columns with 10 ml 1 X PBS (phosphate buffered saline). There  
25 were two 1 m column arranged in tandem each containing the Protein A  
matrix.
- 45 2. Add 3 ml of the serum to 3 ml of PBS and divide this mixture between the  
two columns.
- 30 3. Collect the serum into a test tube as it drains through the column.
- 50 4. When the serum finishes draining, pour the washed serum back into the  
column and begin collecting flow through again. Repeat this step 5 to 6

-19-

times.

5. Wash the columns with 10 ml of 1 X PBS.
6. Prepare several 1 ml tubes with 50  $\mu$ l of 1 M TRIS (2-amino-2-hydroxymethyl-1,3-propanediol) (pH = 9,5).
7. Add 1 ml of elution buffer (100 mM glycine, pH = 2.8) to each tube and collect 1 ml of flow therethrough.
8. Move to the next prepare tube and repeat step 7.
9. Test each 1 ml sample by preparing ELISA plate with 10  $\mu$ l of Bradford Assay and add 50 $\mu$ l of each 1 ml flow through. Keep the samples that change the Bradford Assay from red to blue.
10. Dialyse the positive 1 ml samples together in 4 litres of 1X PBS at pH = 7.2 for at least 24 hours.
11. Use spectrometer at 280 nm to find concentration of IgG in solution (extinction coefficient = 1.4).
12. To store IgG solution, keep frozen at -4°C to -20°C.

Table 5 shows the results for certain antibodies.

Table 5

Peptide used to produce Antibody	Serum used (ml)	Purified Ab volume (ml)	OD <sub>280</sub>	IgG (mg/ml)	Total IgG (mg)
A-1	10	15	3.80	2.71	40.71
A-2	10	15	2.13	1.52	22.82
A-3	10	15	2.93	2.09	31.39
A-4	10	15	3.57	2.55	38.25
A-5	6	12	3.02	2.16	25.88
B-1	10	15	2.64	1.89	28.28
B-2	6	13	4.94	3.53	45.87
B-3	6	13	5.01	3.58	46.52

-20-

Peptide used to produce Antibody	Serum used (ml)	Purified Ab volume (ml)	OD <sub>280</sub>	IgG (mg/ml)	Total IgG (mg)
B-4	10	15	2.68	1.91	28.71
B-5	10	15	2.28	1.63	24.43
B-6	10	15	2.50	1.79	26.78
B-7	10	15	2.90	2.07	31.07
B-8	10	15	3.40	2.43	36.43
B-9	10	15	3.80	2.71	40.71
B-10	10	15	4.18	2.99	44.79
C-12	10	15	1.95	1.39	20.89
D-1	10	15	2.32	1.66	24.86
A-7	6	12	3.33	2.38	28.54

The level of antibodies in the serum was established by ELISA (enzyme-linked immunosorbent assay) with the corresponding synthetic peptide antigen. This technique involved the following steps:

1. The antigen was diluted with a 0.1 M bicarbonate buffer (pH 9.0) to yield a 10 µg of antigen/ml solution. A volume of 50 µl of this solution was placed into each microwell of a 96 well plate.
2. The plates were covered and incubated at 37°C for 3 hours.
3. The wells were washed with a coupling buffer and blocked using 200 µl of Pierce standard solution of BSA (bovine serum albumin).
4. 50 µl of diluent BSA (0.75% soln.) was pipetted into each well. 50 µl of antibody serum sample diluted 1:100 in diluent BSA were placed in lane A of each row.
5. 1:2 serial dilutions were performed moving down the plate.
6. The plates were covered and incubated at room temperature for 60 minutes.
7. The plates were washed four times with PBS wash solution.
8. A volume of 50 µl of goat anti-rabbit IgG (H&L) HRP conjugate at 1:1000

-21-

dilution in BSA was pipetted into each well and incubated at room temperature for 60 minutes (H&L = heavy and light chain; HRP = horseradish peroxidase).

9. The plates were washed four times with PBS wash solution.

10. A volume of 50  $\mu$ l of substrate solution 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid)-diammonium salt (ABTS available from Pierce, which is used to help visualise the extent of the antibody/antigen reaction) was pipetted into each well and incubated at room temperature for about 2 minutes.

11. The reaction was stopped by adding 50  $\mu$ l of 1% SDS (sodium dodecyl sulfate) into each well.

12. The wells were then read on a dynoplate reader at 405.

The data presented in Table 6 show the serum antibody titers against specific antibodies after the 10 week immunisation protocol.

Table 6

<u>No</u>	<u>Sequences</u>	<u>Titre: (Serum Dilution)</u>			
		<u>Pre</u>		<u>Post</u>	
		<u>Immunization</u>		<u>Immunization</u>	
20	<b>A. Peptides of unknown origin</b>	<b>R1</b>	<b>R2</b>	<b>R1</b>	<b>R2</b>
1	LQTPQPLLQVMMEPQGD	0	0	6400	0
2	MPQNFYKLPQM	0	0	6400	25600
3	VLEMKFPPPPQETVT	0	0	6400	12800
4	LKPFPKLKVEVFPFP	0	0	6400	25600
25	5 SEQP	0	0	3200	25600
6	DKE	ND	ND	ND	ND
7	DPPPPQS	0	0	3400	6200
8	LNf	ND	ND	ND	ND
	<b>B. Peptides from casein homologue</b>	<b>R1</b>	<b>R2</b>	<b>R1</b>	<b>R2</b>

5

-22-

10

15

20

25

30

35

40

45

50

55

1	VLPPNVG	0	0	25600	25600
2	KYKLQPE	0	0	25600	25600
3	SEEMP	0	0	25600	12800
4	DSQPPV	0	0	25600	25600
5	5 FPPPK	0	0	12800	6400
6	VVMEV	0	0	25600	25600
7	DLEMPVLPVEPFPFV	0	0	25600	6400
8	LFFFLPVWNVLP	0	0	200	200
9	MQPPPLP	0	0	3200	12800
10	10 DQPPDVEKPDQLQPFQVQS	0	0	12800	25600
<b>C. Peptides from <math>\beta</math>-casein</b>		<b>R1</b>	<b>R2</b>	<b>R1</b>	<b>R2</b>
1	VYPFTGPIPN	ND	0	ND	>10000
2	SLPQNILPL	ND	0	ND	>10000
3	TQTPVVVPPF	ND	0	ND	>10000
4	15 LQPEIMGVPKVKEMVPK	ND	0	ND	>10000
5	HKEMPFPKYPVEPFOTESQ	ND	0	ND	>10000
6	SLTLTDVEKLHLPLPLVQ	ND	0	ND	>10000
7	SWMHQPP	ND	ND	ND	ND
8	QPLPPTVMFP	ND	ND	ND	ND
9	20 MHQPPQPLPPTVMFP	ND	0	ND	>10000
10	PQSVLS	ND	ND	ND	ND
11	LSQPKVLPVPQKAVPQRDMPIQ	ND	0	ND	>10000
12	AFLLYQE	ND	0	12800	25600
13	FLLYQEPVLGPVR	ND	0	ND	>10000
14	25 RGPFPILV	ND	ND	ND	ND
<b>D. Peptide from annexin</b>		<b>R1</b>	<b>R2</b>	<b>R1</b>	<b>R2</b>
1	ATFNRYQDDHGEEILKSL	0	0	12800	25600



ND = Not Done

In Table 6 the results are shown for two rabbits R1 and R2. In general, these results indicate that the potency of the antibodies produced in respect of peptides was excellent, and therefore that each antibody was the correct antibody for its synthetic peptide antigen. The antibodies produced by this technique were monospecific. However, the antigenic response in respect of peptides A-1, A-7 and B-8 were significantly lower than expected and lead us to predict that these peptides, especially B-8, would be useful as an immunosuppressant, and therefore would be useful in the treatment of autoimmune disorder and in the prevention of organ rejection during, for example, organ transplants.

#### Example 4

In order to establish that the peptides corresponding to the synthetic peptide antigens exist in Colostrinin we carried out tests to determine whether certain of the antibodies produced a reaction in Colostrinin itself.

We studied the rate at which the peptides A-4, B-7, B-8 and B-9 disappeared from colostrum produced in sheep. The colostrum was collected from the mother's milk at 24 hours, 48 hours and 72 hours post parturition, and the level of the peptides was measured. The peptide level was measured by means of an antigen-antibody reaction, using the antibodies produced by the method of Example 3. The result are shown in Table 7.

Table 7

Peptide:	24 Hour Titre	48 Hour Titre	72 Hour Titre
A-4	12800	6400	3200
B-7	12800	6400	3200
B-8	12800	3200	3200
B-9	12800	12800	3200

These results demonstrated that antibodies had recognised the amino acid

5

sequences A-4, B-7, B-8 and B-9, and that the concentration of the peptide had diminished over time, owing to binding of the antibody with the peptide.

10

It will be appreciated that the invention described above may be modified.

15

20

25

30

35

40

45

50

55

5

-25-

CLAIMS:

10

5

15

20

25

30

35

40

45

50

1. A peptide, in substantially isolated form, which substantially includes the amino-terminal amino acid sequence: LQTPQPLLQVMMEPQGD-OH (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLVFVFPF (SEQ ID 4); SEQP (SEQ ID 5); DKE (SEQ ID 6); DPPPPQS (SEQ ID 7); LNF (SEQ ID 8); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); SEEMP (SEQ ID 11); DSQPPV (SEQ ID 12); FPPPK (SEQ ID 13); VMEV (SEQ ID 14); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVVNVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS (SEQ ID 18); VYPFTGPIPN (SEQ ID 19); SLPQNILPL (SEQ ID 20); TQTPVVVPF (SEQ ID 21); LQPEIMGVVKVETMVPK (SEQ ID 22); HKEMFPKYPVEPFTESQ (SEQ ID 23); SLTLTDVEKLHLPLPLVQ (SEQ ID 24); SWMHQPP (SEQ ID 25); QPLPPTVMFP (SEQ ID 26); MHQPPQLPPTVMFP (SEQ ID 27); PQSVLS (SEQ ID 28); LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID 29); AFLLYQE (SEQ ID 30); FLLYQEPVLGPVR (SEQ ID 31); RGPFPILV (SEQ ID 32); ATFNRYQDDHGEEILKSL (SEQ ID 33).

2. A peptide, in substantially isolated form, which substantially includes the amino acid sequence: LQTPQPLLQVMMEPQGD (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLVFVFPF (SEQ ID 4); DPPPPQS (SEQ ID 7); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); DSQPPV (SEQ ID 12); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVVNVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS (SEQ ID 18).

3. A peptide, in substantially isolated form, which substantially entirely consists of the amino acid sequence: LQTPQPLLQVMMEPQGD (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLVFVFPF (SEQ ID 4); SEQP (SEQ ID 5); DKE (SEQ ID 6); DPPPPQS (SEQ ID 7); LNF (SEQ ID 8); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); SEEMP (SEQ ID 11); DSQPPV (SEQ ID 12); FPPPK (SEQ ID 13); VMEV (SEQ ID 14); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVVNVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS

55

-26-

(SEQ ID 18); VYPFTGPIPN (SEQ ID 19); SLPQNILPL (SEQ ID 20); TQTPVWVPF (SEQ ID 21); LQPEIMGVPKVKETMVPK (SEQ ID 22); HKEMPFKYPVEPFTESQ (SEQ ID 23); SLTLTDVEKLHLPLPLVQ (SEQ ID 24); SWMHQPP (SEQ ID 25); QPLPPTVMFP (SEQ ID 26); MHQPPQPLPPTVMFP (SEQ ID 27); PQSVLS (SEQ ID 28); LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID 29); AFLLYQE (SEQ ID 30); FLLYQEPVLGPVR (SEQ ID 31); RGPFPILV (SEQ ID 32); ATFNRYQDDHGEEILKSL (SEQ ID 33).

4. A peptide according to claim 1, 2 or 3, when obtained by a synthetic process.

10

5. A peptide obtained by a synthetic process, which substantially includes the amino-terminal amino acid sequence: LQTPQPLLQVMMEPQGD (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLKVEVFPPF (SEQ ID 4); SEQP (SEQ ID 5); DKE (SEQ ID 6); DPPPPQS (SEQ ID 7); LNF (SEQ ID 8); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); SEEMP (SEQ ID 11); DSQPPV (SEQ ID 12); FPPPK (SEQ ID 13); VMEV (SEQ ID 14); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVVNVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS (SEQ ID 18); VYPFTGPIPN (SEQ ID 19); SLPQNILPL (SEQ ID 20); TQTPVWVPF (SEQ ID 21); LQPEIMGVPKVKETMVPK (SEQ ID 22); HKEMPFKYPVEPFTESQ (SEQ ID 23); SLTLTDVEKLHLPLPLVQ (SEQ ID 24); SWMHQPP (SEQ ID 25); QPLPPTVMFP (SEQ ID 26); MHQPPQPLPPTVMFP (SEQ ID 27); PQSVLS (SEQ ID 28); LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID 29); AFLLYQE (SEQ ID 30); FLLYQEPVLGPVR (SEQ ID 31); RGPFPILV (SEQ ID 32); ATFNRYQDDHGEEILKSL (SEQ ID 33).

25

6. A peptide obtained by a synthetic process, which substantially includes the amino acid sequence: LQTPQPLLQVMMEPQGD (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLKVEVFPPF (SEQ ID 4); DPPPPQS (SEQ ID 7); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); DSQPPV (SEQ ID 12); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVVNVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS (SEQ ID 18).

-27-

7. A peptide obtained by a synthetic process, which substantially entirely consists of the amino acid sequence: LQTPQPLLQVMMEPQGD (SEQ ID 1); MPQNFYKLPQM (SEQ ID 2); VLEMKFPPPPQETVT (SEQ ID 3); LKPFPKLKVEVFPPF (SEQ ID 4); SEQP (SEQ ID 5); DKE (SEQ ID 6); DPPPQSQS (SEQ ID 7); LNF (SEQ ID 8); VLPPNVG (SEQ ID 9); KYKLQPE (SEQ ID 10); SEEMP (SEQ ID 11); DSQPPV (SEQ ID 12); FPPPK (SEQ ID 13); VMEV (SEQ ID 14); DLEMPVLPVEFPFV (SEQ ID 15); LFFFLPVWVLP (SEQ ID 16); MQPPPLP (SEQ ID 17); DQPPDVEKPDLPFQVQS (SEQ ID 18); VYPFTGPIPN (SEQ ID 19); SLPQNILPL (SEQ ID 20); TQTPVVPPF (SEQ ID 21); LQPEIMGVVKVETMVPK (SEQ ID 22); HKEMPFKYPVEPFTESEQ (SEQ ID 23); SLTLTDVEKLHLPLVQ (SEQ ID 24); SWMHQPP (SEQ ID 25); QPLPPTVMFP (SEQ ID 26); MHQPPQPLPPTVMFP (SEQ ID 27); PQSVLS (SEQ ID 28); LSQPKVLPVPQKAVPQRDMPIQ (SEQ ID 29); AFLLYQE (SEQ ID 30); FLLYQEPVLGPVR (SEQ ID 31); RGPFPILV (SEQ ID 32); ATFNRYQDDHGEEILKSL (SEQ ID 33).

8. A peptide comprising:  $\text{NH}_2$ -(Ac)CLQTPQPLLQVMMEPQGD-OH (SEQ ID 34);  $\text{NH}_2$ -(Ac)CMPQNFYKLPQM-OH (SEQ ID 35);  $\text{NH}_2$ -(Ac)CVLEMKFPPPPQETVT-OH (SEQ ID 36);  $\text{NH}_2$ -(Ac)CLKPFPKLKVEVFPPF-OH (SEQ ID 37);  $\text{NH}_2$ -SEQPGGGC-OH (SEQ ID 38);  $\text{NH}_2$ -(Ac)CGVLPPNVG-OH (SEQ ID 39);  $\text{NH}_2$ -(Ac)CGGGKYKLQE-OH (SEQ ID 40);  $\text{NH}_2$ -(Ac)CGGGSEEMP(amide)-OH (SEQ ID 41);  $\text{NH}_2$ -(Ac)CGGGDSQPPV-OH (SEQ ID 42);  $\text{NH}_2$ -CFPPPKGGGC-OH (SEQ ID 43);  $\text{NH}_2$ -(Ac)CGGGVMEV-OH (SEQ ID 44);  $\text{NH}_2$ -(Ac)CDLEMPVLPVEFPFV-OH (SEQ ID 45);  $\text{NH}_2$ -(Ac)CLFFFLPVWVLP-OH (SEQ ID 46);  $\text{NH}_2$ -(Ac)CMQPPPLP-OH (SEQ ID 47);  $\text{NH}_2$ -(Ac)CDQPPDVEKPDLPFQVQS-OH (SEQ ID 48);  $\text{NH}_2$ -(Ac)CGAFLLYQE-OH (SEQ ID 49);  $\text{NH}_2$ -(Ac)CATFNRYQDDHGEEILKSL-OH (SEQ ID 50).

9. A peptide according to any preceding claim, for use as a medicament.

10. A peptide according to claim 9, for use in the treatment of chronic disorders of the central nervous system.

-28-

11. A peptide according to claim 10, for use in the treatment of neurological disorders and/or mental disorders.

12. A peptide according to claim 9, for use in the treatment of dementia and/or neurodegenerative diseases.

13. A peptide according to claim 9, for use in the treatment of Alzheimer's disease and/or motor neurone disease.

14. A peptide according to claim 9, for use in the treatment of psychosis and/or neurosis.

15. A peptide according to claim 9, for use in the treatment of chronic disorders of the immune system.

16. A peptide according to claim 9, for use in the treatment of diseases with a bacterial and viral aetiology, and/or for use in the treatment of acquired immunological deficiencies.

17. A peptide according to claim 9, for use in the treatment of chronic bacterial and/or viral infections.

18. A peptide according to claim 9, for use in the treatment of diseases characterised by the presence of  $\beta$ -amyloid plaque.

19. The use of a peptide according to any one of claims 1 to 8, in the manufacture of a medicament for the treatment of chronic disorders of the central nervous system.

20. The use of a peptide according to any one of claims 1 to 8 in the manufacture of a medicament for the treatment of chronic disorders of the immune system.

5

-29-

10

21. A method of treating disorders of the central nervous system and/or of the immune system, comprising administering a therapeutically effective amount of a peptide according to any one of claims 1 to 8 to a patient.

15

22. A composition comprising a peptide according to any one of claims 1 to 8, in combination with a physiologically acceptable carrier.

20

23. A composition comprising two or more peptides according to any one of claims 1 to 8, in combination with a physiologically acceptable carrier.

10

24. A composition according to claim 22 or 23, in a form suitable for injection.

25

25. A composition according to claim 22 or 23, in a form suitable for absorption through the mucosa of the oral/nasopharyngeal cavity and/or in a form suitable for absorption in the alimentary canal.

30

26. A composition according to claim 22 or 23, in the form of a tablet, lozenge, gel, patch or plaster.

35

27. A composition according to claim 22 or 23, in a form suitable for topical application.

40

28. The use of a peptide according to any one of claims 1 to 8 as a dietary supplement.

25

45

29. The use of a peptide according to any one of claims 1 to 8 as a dietary supplement for babies, small children, adults who have been subjected to chemotherapy and/or adults who have suffered from anorexia, or weight loss due to chronic disease.

50

30. A dietary supplement comprising an orally ingestible combination of a peptide according to any one of claims 1 to 8 combination with a physiologically acceptable

55

5

-30-

carrier.

10

31. An antibody which binds to a peptide according to any one of claims 1 or 8.

5 32. An antibody obtainable by using a peptide according to any one of claims 1 to 8 as an antigen.

15

33. A peptide containing the amino acid sequence LQTPQPLLQVMMEPQGD; DPPPQS; and/or LFFFLPVNVLP for use as an immunosuppressant.

20

10

34. A peptide containing the amino acid sequence LQTPQPLLQVMMEPQGD; DPPPQS; and/or LFFFLPVNVLP for use in the treatment of autoimmune disorder.

25

35. A peptide containing the amino acid sequence LQTPQPLLQVMMEPQGD; DPPPQS; and/or LFFFLPVNVLP for use in suppressing the rejection of transplanting organs.

15

30

35


40


45


50


55













5/18




SUBSTITUTE SHEET (RULE 26)


6/18



7/18




SUBSTITUTE SHEET (RULE 26)






9/18




10/18




SUBSTITUTE SHEET (RULE 26)


11/18




12/18




SUBSTITUTE SHEET (RULE 26)



14/18




15/18



SUBSTITUTE SHEET (RULE 26)


16/18




SUBSTITUTE SHEET (RULE 26)



17/18



SUBSTITUTE SHEET (RULE 26)



-1-

## SEQUENCE LISTING

&lt;110&gt; ReGen Therapeutics plc

&lt;120&gt; Peptides

&lt;130&gt; PAC/19599

&lt;140&gt;

&lt;141&gt;

&lt;150&gt; 9912852.2

&lt;151&gt; 1999-06-02

&lt;160&gt; 50

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 17

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 1

Leu	Gln	Thr	Pro	Gln	Pro	Leu	Leu	Gln	Val	Met	Met	Glu	Pro	Gln	Gly
1				5					10					15	

Asp

&lt;210&gt; 2

&lt;211&gt; 11

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 2

Met	Pro	Gln	Asn	Phe	Tyr	Lys	Leu	Pro	Gln	Met
1				5					10	

&lt;210&gt; 3

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

-2-

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 3

Val Leu Glu Met Lys Phe Pro Pro Pro Pro Gln Glu Thr Val Thr  
1 5 10 15

<210> 4

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 4

Leu Lys Pro Phe Pro Lys Leu Lys Val Glu Val Phe Pro Phe Pro  
1 5 10 15

<210> 5

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 5

Ser Glu Gln Pro  
1

<210> 6

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 6

Asp Lys Glu  
1

<210> 7

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

-3-

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 7

Asp Pro Pro Pro Pro Gln Ser  
1 5

<210> 8

<211> 3

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 8

Leu Asn Phe  
1

<210> 9

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 9

Val Leu Pro Pro Asn Val Gly  
1 5

<210> 10

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 10

Lys Tyr Lys Leu Gln Pro Glu  
1 5

<210> 11

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

-4-

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 11  
Ser Glu Glu Met Pro  
1 5

<210> 12  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 12  
Asp Ser Gln Pro Pro Val  
1 5

<210> 13  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 13  
Phe Pro Pro Pro Lys  
1 5

<210> 14  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 14  
Val Val Met Glu Val  
1 5

<210> 15  
<211> 15  
<212> PRT  
<213> Artificial Sequence

<220>

-5-

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 15  
Asp Leu Glu Met Pro Val Leu Pro Val Glu Pro Phe Pro Phe Val  
1 5 10 15

<210> 16

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 16  
Leu Phe Phe Phe Leu Pro Val Val Asn Val Leu Pro  
1 5 10

<210> 17

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 17  
Met Gln Pro Pro Pro Leu Pro  
1 5

<210> 18

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 18  
Asp Gln Pro Pro Asp Val Glu Lys Pro Asp Leu Gln Pro Phe Gln Val  
1 5 10 15

Gln Ser

<210> 19

<211> 10

<212> PRT

SUBSTITUTE SHEET (RULE 26)

- 6 -

## &lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 19

Val Tyr Pro Phe Thr Gly Pro Ile Pro Asn  
1 5 10

&lt;210&gt; 20

&lt;211&gt; 9

&lt;212&gt; PRT

## &lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 20

Ser Leu Pro Gln Asn Ile Leu Pro Leu  
1 5

&lt;210&gt; 21

&lt;211&gt; 10

&lt;212&gt; PRT

## &lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 21

Thr Gln Thr Pro Val Val Val Pro Pro Phe  
1 5 10

&lt;210&gt; 22

&lt;211&gt; 18

&lt;212&gt; PRT

## &lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 22

Leu Gln Pro Glu Ile Met Gly Val Pro Lys Val Lys Glu Thr Met Val  
1 5 10 15

Pro Lys



-7-

<210> 23  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 23  
His Lys Glu Met Pro Phe Pro Lys Tyr Pro Val Glu Pro Phe Thr Glu  
1 5 10 15

Ser Gln

<210> 24  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 24  
Ser Leu Thr Leu Thr Asp Val Glu Lys Leu His Leu Pro Leu Pro Leu  
1 5 10 15

Val Gln

<210> 25  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 25  
Ser Trp Met His Gln Pro Pro  
1 5

<210> 26  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

-8-

&lt;400&gt; 26

Gln Pro Leu Pro Pro Thr Thr Val Met Phe Pro  
1 5 10

&lt;210&gt; 27

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 27

Met His Gln Pro Pro Gln Pro Leu Pro Pro Thr Val Met Phe Pro  
1 5 10 15

&lt;210&gt; 28

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 28

Pro Gln Ser Val Leu Ser  
1 5

&lt;210&gt; 29

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 29

Leu Ser Gln Pro Lys Val Leu Pro Val Pro Gln Lys Ala Val Pro Gln  
1 5 10 15

Arg Asp Met Pro Ile Gln  
20

&lt;210&gt; 30

&lt;211&gt; 7

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

-9-

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 30  
Ala Phe Leu Leu Tyr Gln Glu  
1 5

<210> 31  
<211> 13  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 31  
Phe Leu Leu Tyr Gln Glu Pro Val Leu Gly Pro Val Arg  
1 5 10

<210> 32  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 32  
Arg Gly Pro Phe Pro Ile Leu Val  
1 5

<210> 33  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 33  
Ala Thr Phe Asn Arg Tyr Gln Asp Asp His Gly Glu Glu Ile Leu Lys  
1 5 10 15

Ser Leu

<210> 34  
<211> 18  
<212> PRT

-10-

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 34

Cys	Leu	Gln	Thr	Pro	Gln	Pro	Leu	Leu	Gln	Val	Met	Met	Glu	Pro	Gln
1				5				10					15		

Gly Asp

&lt;210&gt; 35

&lt;211&gt; 12

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 35

Cys	Met	Pro	Gln	Asn	Phe	Tyr	Lys	Leu	Pro	Gln	Met
1				5				10			

&lt;210&gt; 36

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 36

Cys	Val	Leu	Glu	Met	Lys	Phe	Pro	Pro	Pro	Pro	Gln	Glu	Thr	Val	Thr
1				5					10				15		

&lt;210&gt; 37

&lt;211&gt; 16

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 37

Cys	Leu	Lys	Pro	Phe	Pro	Lys	Leu	Lys	Val	Glu	Val	Phe	Pro	Phe	Pro
1				5				10					15		

-11-

<210> 38  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 38  
Ser Glu Gln Pro Gly Gly Gly Cys  
1 5

<210> 39  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 39  
Cys Gly Val Leu Pro Pro Asn Val Gly  
1 5

<210> 40  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 40  
Cys Gly Gly Gly Lys Tyr Lys Leu Gln Glu  
1 5 10

<210> 41  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 41  
Cys Gly Gly Gly Ser Glu Glu Met Pro  
1 5

-12-

<210> 42  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 42  
Cys Gly Gly Gly Asp Ser Gln Pro Pro Val  
1 5 10

<210> 43  
<211> 10  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 43  
Cys Phe Pro Pro Pro Lys Gly Gly Gly Cys  
1 5 10

<210> 44  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 44  
Cys Gly Gly Gly Val Val Met Glu Val  
1 5

<210> 45  
<211> 16  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 45  
Cys Asp Leu Glu Met Pro Val Leu Pro Val Glu Pro Phe Pro Phe Val  
1 5 10 15

-13-

<210> 46  
<211> 14  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 46  
Cys Leu Phe Phe Phe Leu Pro Val Val Asn Val Leu Pro Ile  
1 5 10

<210> 47  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 47  
Cys Met Gln Pro Pro Pro Leu Pro  
1 5

<210> 48  
<211> 19  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 48  
Cys Asp Gln Pro Pro Asp Val Glu Lys Pro Asp Leu Gln Pro Phe Gln  
1 5 10 15  
Val Gln Ser

<210> 49  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

<400> 49  
Cys Gly Ala Phe Leu Leu Tyr Gln Glu

-14-

1

5

&lt;210&gt; 50

&lt;211&gt; 19

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Description of Artificial Sequence: Peptide  
isolated from colostrinin

&lt;400&gt; 50

Cys Ala Thr Phe Asn Arg Tyr Gln Asp Asp His Gly Glu Glu Ile Leu  
1 5 10 15

Lys Ser Leu